

## FITC AffiniPure Goat Anti-Rabbit IgG(H+L)



<u>Catalog No.</u>	<u>Size</u>
E031220-01	100µl
E031220-02	500µl

<b>Product Type</b>	Secondary Antibody
<b>Physical State</b>	Liquid
<b>Antibody Source</b>	The antibody was purified from antisera by immunoaffinity chromatography using antigens coupled to agarose beads
<b>Antibody Purity</b>	The antibody reacts with whole molecule rabbit IgG. It also reacts with the light chains of other rabbit immunoglobulins.
<b>Suggested Dilution Range</b>	1:50 - 1:200 for most applications
<b>Fluorophore</b>	Fluorescein-5-isothiocyanate (FITC-isomer 1) Amax=492 nm; Emax=520 nm
<b>Fluorophore/Protein Ratio</b>	8.2µg/mg; 3.1moles FITC per mole IgG
<b>Storage Buffer</b>	0.01M Sodium phosphate, 0.25M NaCl, 50% Glycerol, pH7.6
<b>Stabilizer</b>	15 mg/ml Bovine Serum Albumin (IgG-Free, Protease-Free)
<b>Preservative</b>	0.05% Sodium Azide
<b>Storage Instructions</b>	Store at -20°C. Avoid repeated freeze / thaw cycles.
<b>Note</b>	The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Nothing disclosed herein is to be construed as a recommendation to use this product in violation of any patents. The information presented above is believed to be accurate. However, said information and product are offered without warranty or guarantee since the ultimate conditions of use and the variability of the materials treated are beyond our control. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.